



# CENOL 0.5%

## MULTIPURPOSE INSECTICIDE

**FOR USE ONLY BY PEST CONTROL OPERATORS  
OR BY PUBLIC HEALTH OFFICIALS  
AND TRAINED PERSONNEL OF  
MOSQUITO ABATEMENT DISTRICTS  
AND OTHER MOSQUITO CONTROL PROGRAMS**

• A READY-TO-USE PERMETHRIN STABLE EMULSION / INSECTICIDE  
FOR USE OUTDOORS, IN HOMES AND NON-FOOD AREAS OF  
FOOD/FEED PROCESSING ESTABLISHMENTS, KENNELS, COMMERCIAL  
BUILDINGS, WAREHOUSES, THEATERS, OFFICE BUILDINGS,  
SCHOOLS, MOTELS, AND HOTELS

• FOR CONTROL OF FLEAS, BROWN DOG TICKS AND LICE  
ON PREMISES AND ON DOGS

• FOR CONTROL OF TICKS WHICH MAY CAUSE LYME  
DISEASE ON PREMISES

• CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS

• FOR CONTROL OF PESTS ON LIVESTOCK

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

Harmful if swallowed.

See Rear panel for Additional Precautions.

EPA Est. No. 45385-OH-01

EPA Reg. No. 45385-99

### ACTIVE INGREDIENTS:

Permethrin..... 0.50%

INERT INGREDIENT ..... 99.50%

**TOTAL 100.00%**

(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl)  
2,2-dimethylcyclopropanecarboxylate.  
Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.

**SHAKE WELL BEFORE USING**

**READ ENTIRE LABEL BEFORE EACH USE  
PET USE APPLIES TO USE ONLY ON DOGS**

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before space spraying or surface applications. Do not use on humans. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark. Do not contaminate water when disposing of equipment washwaters.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

### FIRST AID

<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>

### DIRECTIONS FOR USE

It is a violation of the Federal Law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **Pesticide Storage And Spill Procedures:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

**Pesticide Disposal:** Pesticide or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

#### Container Disposal:

**CONTAINERS ONE GALLON AND SMALLER:** Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

**CONTAINERS LARGER THAN ONE GALLON:**

**Metal Containers-** Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

**Plastic Containers -** Triple rinse or equivalent, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Glass Containers-** Triple rinse or equivalent, then dispose of in a sanitary landfill or by other approved state and local procedures.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at **1-800-858-7378**



**CENOL**  
INCORPORATED  
P.O. Box 472  
Twinsburg, OH 44087

See Other Side

**Net Contents:** ■ One Gallon ■ Five Gallons  
■ 30 Gallons ■ 55 Gallons

# SPECIMEN LABEL

## INDOOR APPLICATION

Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed.

Nonfood/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Vacate treated areas and ventilate before reoccupying. For maximum effectiveness, a combination of spot, surface treatment and fogging is recommended.

Do not apply to classrooms while in use.

## SURFACE SPRAYING:

**TO CONTROL COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, MILLIPEDES, CENTIPEDES, SOWBUGS, PILLBUGS, ANTS, SILVERFISH, FIREBRATS, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES AND CADELLES:** Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry.

**TO CONTROL CARPET BEETLES:** Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices, and infested areas of shelving. Repeat treatment as needed.

**TO CONTROL BED BUGS:** Spray mattresses lightly particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floor. Repeat treatment as needed. Allow all treated articles to dry thoroughly before use.

**TO CONTROL FLEAS (ADULT AND LARVAE) AND TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (ADULTS AND LARVAE)** Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

**TO CONTROL CLOTHES MOTHS:** Remove any infested articles from storage, brush thoroughly, and air for several hours in sun light, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining. Allow all treated articles to dry thoroughly before use.

**TO KILL LICE AND LOUSE EGGS:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

## SPACE SPRAYING:

Close all doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of **House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Bees, Black Flies, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths and Tobacco Moths:** Direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for 2 hours. Ventilate before reoccupying.

## FOGGING:

To kill exposed or accessible stages of the above named crawling insects use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly when possible. Keep area closed for 2 hours. Ventilate before reoccupying.

## ON LIVESTOCK:

To protect cattle (beef and dairy), goats, sheep, hogs, and horses from Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats, apply a light mist sufficient to wet the surface of the hair.

For face flies, spray face and head, but do not spray into eyes.

## TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES:

Apply at rate of 2 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

**TO CONTROL BLOOD-SUCKING LICE:** Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

**TO CONTROL POULTRY LICE:** Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

**TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES:** Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

## ON DOGS:

To control Fleas, Ticks, and Lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Use protective gloves or mitts when applying product. Repeat application every two weeks, if necessary. **DO NOT USE ON CATS OR KITTENS.**

Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Do not use on dogs under 12 weeks.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

## OUTDOOR APPLICATION

**TO CONTROL TICKS, INCLUDING TICKS WHICH MAY CARRY LYME DISEASE (PERIMETER TREATMENT):** Use Cenol 0.5% Multipurpose Insecticide as a residual treatment to control Ticks including the **Deer Tick (Ixodes dammini), Lone Star Tick, (Amblyomma americanum), Brown Dog Tick (Rhipicephalus sanguineus), American Dog Tick (Demacantor variabilis) and Western Black-legged Tick (Ixodes pacificus).**

For each areas listed below, apply until surface is wet (approximately 1 gal / 750-1000 ft<sup>2</sup>).

Treat around windows and doors, porches, screens, eaves, patios, garages, under stairways, and in crawl spaces where these pests may enter.

Treat a band of vegetation and leaf litter 6 to 10 feet wide around the outside of buildings to help control ticks occurring there.

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

## FOR CONTROL OF GARDEN INSECTS:

For use on roses, dahlias, asters and other ornamentals to kill **Japanese Beetles** hit by spray as well as **White Flies, Aphids, Exposed Thrips, Spiders, Red Mites and Leafminers.** May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

**EFFECTIVE WHEN USED AS DIRECTED AGAINST WHITEFLIES, APHIDS, SPIDER AND RED MITES, EXPOSED THRIPS, LACE BUGS, JAPANESE BEETLES, PAVEMENT ANTS, CLOVER MITES AND ARMYWORMS:** When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

**ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour but do not apply when wind speeds exceed 10 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest. Apply at a rate of 2.5 pints per acre (0.9 ounce per 1000 square feet). Using truck-mounted conventional misting equipment such as Rotomist models 51C, 91 or 100 HT traveling at 5 mph to treat a 300 foot swath, calibrate equipment to deliver 0.95 gallon per minute. Under no conditions should open water be treated with this product. Not to be used within 100 feet (30 meters) of lakes and streams.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

# SPECIMEN LABEL



9 / 2000

EPA Est. No. 45385-OH-01  
EPA Reg. No. 45385-99

ORDER CODE W-650

Page 2 of 2

2 of 2